

# The Open Access Israeli Journal of Aquaculture – Bamidgeh

As from **January 2010** The Israeli Journal of Aquaculture - Bamidgeh (IJA) will be published exclusively as **an on-line Open Access (OA)** quarterly accessible by all AquacultureHub (<http://www.aquaculturehub.org>) members and registered individuals and institutions. Please visit our website (<http://siamb.org.il>) for free registration form, further information and instructions.

This transformation from a subscription printed version to an on-line OA journal, aims at supporting the concept that scientific peer-reviewed publications should be made available to all, including those with limited resources. The OA IJA does not enforce author or subscription fees and will endeavor to obtain alternative sources of income to support this policy for as long as possible.

## Editor-in-Chief

Dan Mires

## Editorial Board

<b>Rina Chakrabarti</b>	Aqua Research Lab, Dept. of Zoology, University of Delhi, India
<b>Angelo Colorni</b>	National Center for Mariculture, IOLR Eilat, Israel
<b>Daniel Golani</b>	The Hebrew University of Jerusalem Jerusalem, Israel
<b>Hillel Gordin</b>	Kibbutz Yotveta, Arava, Israel
<b>Sheenan Harpaz</b>	Agricultural Research Organization Beit Dagan,
<b>Gideon Hulata</b>	Agricultural Research Organization Beit Dagan,
<b>George Wm. Kissil</b>	National Center for Mariculture, IOLR, Eilat, Israel
<b>Ingrid Lupatsch</b>	Swansea University, Singleton Park, Swansea, UK
<b>Spencer Malecha</b>	Dept. of Human Nutrition, Food & Animal Sciences, CTAHR, University of Hawaii
<b>Constantinos Mylonas</b>	Hellenic Center for Marine Research, Crete, Greece
<b>Amos Tandler</b>	National Center for Mariculture, IOLR Eilat, Israel
<b>Emilio Tibaldi</b>	Udine University Udine, Italy
<b>Jaap van Rijn</b>	Faculty of Agriculture, The Hebrew University of Jerusalem, Israel
<b>Zvi Yaron</b>	Dept. of Zoology, Tel Aviv University, Tel Aviv, Israel

Published under auspices of  
**The Society of Israeli Aquaculture and  
Marine Biotechnology (SIAMB),  
University of Hawai'i at Mānoa Library**

&

**University of Hawai'i at Mānoa  
Aquaculture Program**  
in association with  
**AquacultureHub**

<http://www.aquaculturehub.org>



UNIVERSITY  
of HAWAII  
MĀNOA  
LIBRARY



**AquacultureHub**  
educate • learn • share • engage

ISSN 0792 - 156X

© Israeli Journal of Aquaculture - BAMIGDEH.

PUBLISHER:

Israeli Journal of Aquaculture - BAMIGDEH -  
Kibbutz Ein Hamifratz, Mobile Post 25210,  
ISRAEL


Phone: + 972 52 3965809

<http://siamb.org.il>

Copy Editor **Ellen Rosenberg**



The IJA appears exclusively as a peer-reviewed on-line open-access journal at <http://www.siamb.org.il>. To read papers free of charge, please register online at [registration form](#).  
Sale of IJA papers is strictly forbidden.



## Effect of Replacing Live Food with Formulated Feed on Reproductive Performance of Freshwater Angelfish, *Pterophyllum scalare* (Shcultze, 1823)

\*G.S. Shelar<sup>1</sup>, H. Singh<sup>1</sup>, M.M. Shirdhankar<sup>2</sup> and P. Shelar<sup>3</sup>

<sup>1</sup> Department of Aquaculture College of Fisheries, Dr. Balashaeb Sawant Konkan Krishi Vidyapeeth, Ratnagiri - 415 629, Maharashtra, India

<sup>2</sup> Department of Fisheries resources, Economics, Statistics and Extension education, College of Fisheries, Dr. Balashaeb Sawant Konkan Krishi Vidyapeeth, Ratnagiri - 415 629, Maharashtra, India.

<sup>3</sup> Department of Fisheries Environment, Fisheries College and Research Institute, TANVASU, Tuticorin, Tamilnadu, India – 628008

(Received 23.8.13, Accepted 9.10.13)

Key words: *Pterophyllum scalare*, live food, formulated feed, fatty acids, reproductive performance

### Abstract

The combined effect of live food (LF) and formulated feed (FF) on reproductive performance of *Pterophyllum scalare* was studied. Mosquito larvae were used as live food in addition to formulated feed which contained 50% protein and 12% lipid. The experiment was arranged as a completely randomized design (CRD) with six replicates for each treatment: *P. scalare* were fed with formulated feed and/or live food: T1 (100% FF), T2 (75% FF and 25% LF), T3 (50% FF and 50% LF), T4 (25% FF and 75% LF) and T5 (100% LF). Relative fecundity, egg and oil globule diameter was significantly higher ( $P < 0.05$ ) in T4 and T5 diets whereas inter spawning interval was lower in T3, T4, and T5 diets. Similarly hatching percentages were significantly higher with diets T3, T4, and T5. There were no significant differences observed between treatments on relative hatchling number, hatchling length, and weight. No significant difference was observed for egg and whole body composition of *P. scalare* fed with different diets. However there was a significant difference between treatments on fatty acid profile. It was concluded that live food can be replaced by up to 25-50% formulated feed without compromising reproductive performance of *P. scalare*.

## Introduction

Ornamental fish culture is one of the most popular hobbies in developed countries of the world and is also gaining popularity in many developing markets of the world. The growing interest in aquarium fish has resulted in a steady increase in aquarium fish trade globally. According to FAO (2010), export earnings from the ornamental fish trade in US was worth \$1 billion annually and developing countries produced more than 60%. Freshwater cichlids are popular ornamental fish and most of the species are traded worldwide.

The main dietary function is to provide essential nutrients and energy for maintenance of physiological functions, growth, and gonadal development in fish. Most fish breeders use live food to induce breeding in *P. scalare*. It is believed that live food, (which includes worms, insects, and their larvae), induce fish to breed. However live food is not always available. The biggest problem with live food is the possible introduction of diseases into the holding facilities. Comparisons have been made between live food and artificial feed as broodstock diets for *P. scalare* (Degani and Yehuda, 1996), *Xiphophorus helleri* (Kruger et al., 2001; James and Sampath, 2004), *Betta splendens* (James and Sampath, 2002; Mandal et al., 2010) and zebrafish (Markovich et al., 2007). No one has replaced live food with specifically formulated feeds for *P. scalare*. Thus, the aim of the present study was evaluate the effect of partial or total replacement of live feed by formulated feed, on the reproductive performance of *P. scalare*.

## Materials and Methods

Eight to nine month old males and females of *P. scalare* (50 fish) weighing  $16.43 \pm 0.72$  g were reared in captivity in the wet laboratory of Department of Aquaculture, College of Fisheries, Ratnagiri, Maharashtra, India. Prior to initiation of experiments, fish were acclimated to the experimental diets for up to four weeks and were allowed to choose their own mates in a community set-up. A pair was identified in a community tank (180x60x80 cm) when it isolated itself in a corner and drove away other fish in the tank; these pairs were transferred to small spawning tanks (38x23x23 cm) and the experiments were initiated after three spawning cycles of each pair.

Mosquito larvae were used as live food (LF). The formulated feed (FF) was given in different combinations with live feed (LF): T1 (100% FF); T2 (75% FF + 25% LF); T3 (50% FF + 50% LF); T4 (25% FF + 75% LF) and T5 (100% LF). The proximate composition of the experimental diets is presented in Table 1 and fatty acid profile is given in Table 2.

Table 1. Proximate composition (g/kg dry weight basis) of formulated feed and mosquito larvae.

Particulars (g/kg DM)	Formulated feed	Mosquito larvae
Dry matter	964.9	944.3
Crude protein	518.9	498.8
Crude lipid	121.0	141.5
Ash	63.0	173.5
Crude fiber	58	37.2
NFE*	297.1	39.0
Gross energy (KJ/g)**	22.44	16.83

\*NFE = 1000 – (Crude protein + Crude lipid + Ash)

\*\*Gross energy, calculated based on 23.9, 39.8 and 17.6 MJ/kg for protein, lipid and NFE, respectively (Schulz et al., 2007)

Table 2. Fatty acid composition (% of total fatty acids) of formulated feeds and mosquito larvae.

Fatty acid	Formulated feed	Mosquito larvae
<b>Saturates</b>		
12:0	0.4	4.88
14:0	2.7	8.15
16:0	16.9	14.47
18:0	3.4	11.42
20:0	0.4	4.35
22:0	1.5	3.22
Total	25.3	46.49
<b>Monosaturates</b>		
16:1n-7	4.1	-
18:1n-9	33.4	17.78
Total	37.5	17.78
<b>n-6 Polysaturates</b>		
18:2n-6	26.8	6.54
Total	26.8	6.54
<b>n-3 Polysaturates</b>		
18:3n-3	1.8	5.66
20:5n-3	3	3.85
22:6n-3	0.3	1.35
Total	5.1	10.86
Total	31.9	17.4
<b>polyunsaturates</b>		
P:S <sup>1</sup>	1.26	0.37
M:S <sup>2</sup>	1.48	0.38
n-3:n-6	0.19	1.66

<sup>1</sup>P:S = Polyunsaturates : saturates.

<sup>2</sup>M:S = Monounsaturates : saturates.

The experiment was arranged in a completely randomized design (CRD) with six replicates for each treatment. Uneaten feed together with about 40% of the water was siphoned out from each container daily. The hatching rate (%), relative fecundity, was measured as the number of hatchlings (n) in relation to the weight of the parents ( $n/g^{-1}$ ), inter spawning interval (ISI), relative hatchling number, length and weight of free swimming larvae was recorded after each spawning until the fifth consecutive spawn of each pair. On the sixth spawn, the egg and oil globule diameter were estimated by tpsDIG2 software (Rohlf, 2004). The remaining eggs were used for biochemical analysis. At the end of the experiment the broodfish were sacrificed for muscle tissue biochemical analysis.

Each tank (38x23x23 cm) was provided with a spawning slate (11"x3") that was placed vertically along the wall of the aquarium. After spawning, the eggs were counted, the spawning slate was transferred to a vigorously aerated hatching tank (45x22x30 cm), and one ppm methylene blue was added to prevent the growth of fungi. After 3-4 days the hatchlings reached wriggler-stage and remained attached to the spawning slate. After 6-7 days, the hatchlings consumed their yolk sac and became free swimming. These hatchlings were used to evaluate the hatching performance of the fish.

The proximate analyses of feed ingredients, diets, muscle tissue, and eggs, were carried out according to the standard methods of AOAC (2006). Moisture content was estimated by drying the samples to constant weight at 110°C in a drying oven (BTI); protein was determined using the Kjeldahl method (Kel Plus-Classic DX, Pelican, India); and crude fat was determined by ether extraction using Soxhlet apparatus (Socs Plus-SCS 2, Pelican, India). Crude fiber was estimated through 1.25% acid subsequent 1.25% alkali digestion using Fibra plus-FES 2, Pelican, India. Ash was determined by combusting fat free samples in a muffle furnace at 550°C for 5 hours. The egg protein and lipid content was analyzed using Lowry (Lowry et al., 1951) and Folch's (Folch et al., 1957) methods respectively.

**Spawning and hatching performance.** The reproductive parameters such as relative fecundity (RF), time interval between each spawning in days, i.e. inter spawning interval (ISI), as well as egg and oil globule diameter were calculated. The mean of the short and long axes diameters were taken as the diameter of the egg (Gunasekera et al., 1995). Egg and oil globule diameter was measured by tpsDIG2 software (Rohlf, 2004).

$$\text{a. Relative fecundity (nos.)} = \frac{\text{Total Number of eggs produced}}{\text{Total body weight of female (g)}}$$

The weight of hatchling was estimated using the Top pan electronic precision (0.01 mg accuracy) balance. The length of hatchling was recorded using a measuring scale (0.5 mm fraction). Hatchlings were counted and shifted to another rearing tank. The healthy hatchlings were used to calculate relative hatchling number using formulae as follows,

$$\begin{aligned} \text{a. Hatching percent (\%)} &= \frac{\text{Total number of eggs hatched}}{\text{Total number of eggs produced}} \times 100 \\ \text{b. Relative hatchling number} &= \frac{\text{Total number of hatchling produced}}{\text{Total body weight of female (g)}} \times 100 \end{aligned}$$

**Statistical analysis.** Experiments were analyzed by analysis of variance (ANOVA) using Statistical Analysis System (SAS 9.2) and difference among means were tested using Tukey's HSD test (Snedecor and Cochran, 1967). Percent values were transformed into arcsines (Zar, 2005).

## Results

**Spawning and hatching performance of *P. scalare* fed with different type of feeds.** The observed effect of varying levels of formulated feed and live food on average values for relative fecundity, inter spawning interval, egg and oil globule diameter of *P. scalare* is presented in Table 3. The maximum relative fecundity of  $59.52 \pm 0.36$  nos. $g^{-1}$  was

observed in T5 diet which was significantly different ( $P < 0.05$ ) from T1, T2 and T3 diets. Minimum average spawning interval of  $12.12 \pm 0.23$  days was recorded in T4 diet while with the T1 diet, maximum average was  $14.70 \pm 0.36$  days. Significantly higher egg ( $1.3602 \pm 0.0069$  mm) and oil globule diameter ( $1.2243 \pm 0.0015$  mm) was observed in T5 diet.

Average hatching percent, relative hatchling number, length and weight of *P. scalare* observed due to effect of varying levels of formulated feed and live food is presented in Table 3.0.

Table 3.0 Spawning and hatching performance of *P. scalare* fed with different feed types.

Particulars	T1	T2	T3	T4	T5
Relative fecundity (nos. g <sup>-1</sup> )	$55.89 \pm 0.85^b$	$54.50 \pm 1.24^b$	$56.60 \pm 0.63^b$	$59.10 \pm 0.54^a$	$59.52 \pm 0.36^a$
Inter spawning interval (days)	$14.70 \pm 0.36^a$	$14.57 \pm 0.18^a$	$12.72 \pm 0.20^b$	$12.12 \pm 0.23^b$	$12.17 \pm 0.19^b$
Egg diameter (mm)	$1.3262 \pm 0.0073^{ab}$	$1.3233 \pm 0.0075^{ab}$	$1.3209 \pm 0.0055^{ab}$	$1.3426 \pm 0.0022^a$	$1.3602 \pm 0.0069^a$
Oil globule diameter (mm)	$1.2003 \pm 0.0037^b$	$1.2020 \pm 0.0026^b$	$1.2016 \pm 0.0021^b$	$1.2193 \pm 0.0035^a$	$1.2243 \pm 0.0015^a$
Hatching percent %	$68.08 \pm 0.98^b$	$67.43 \pm 0.80^b$	$70.18 \pm 0.40^a$	$70.90 \pm 0.32^a$	$71.75 \pm 0.67^a$
Relative hatchling number	$41.46 \pm 0.21$	$41.21 \pm 0.57$	$41.57 \pm 0.22$	$41.57 \pm 0.23$	$41.92 \pm 0.56$
Hatchling length (cm)	$0.52 \pm 0.02$	$0.50 \pm 0.00$	$0.52 \pm 0.02$	$0.53 \pm 0.02$	$0.55 \pm 0.02$
Hatchling weight (mg)	$4.35 \pm 0.08$	$4.33 \pm 0.11$	$4.48 \pm 0.08$	$4.57 \pm 0.07$	$4.50 \pm 0.06$

Mean values in similar row with different letters are significantly different (Tukey's HSD test,  $P < 0.05$ ).

T<sub>1</sub>=100% FF; T<sub>2</sub>=75% FF + 25% LF; T<sub>3</sub>=50% FF + 50% LF; T<sub>4</sub>=25% FF + 75% LF; T<sub>5</sub>=100% LF.

Significantly maximum hatching percent of  $71.75 \pm 0.67$  % was observed in T5 diet whereas the minimum of  $67.43 \pm 0.80$  % was seen in the T2 diet. This was higher than in diets T1 and T2. The maximum relative hatchling number  $41.92 \pm 0.56$  was observed in T5 diet and the lowest  $41.21 \pm 0.57$  was seen in T2. Maximum average hatchling length ( $0.55 \pm 0.02$  cm) and weight ( $4.57 \pm 0.07$  mg) was recorded in T5 and T4 diets respectively. There was no significant difference for relative hatchling number, hatchling length and weight.

**Biochemical composition of eggs and muscle tissue of *P. scalare*.** The average moisture, protein and lipid content of eggs of *P. scalare* fed with different levels of formulated feed and live food is given in Table 4. ANOVA test did not show a significant difference ( $P > 0.05$ ) between moisture, protein, and lipid percent of eggs.

The average values for moisture, protein, lipid, and ash content of muscle tissue of *P. scalare* fed with different levels of formulated feed and live food are given in Table 4. There was no significant difference ( $P > 0.05$ ) for muscle moisture, protein, lipid, and ash between the different treatments.

Table 4. Proximate composition of eggs (% wet weight basis) and muscle composition (% dry weight basis) of *P. scalare* fed with different feed type.

Particulars	T1	T2	T3	T4	T5
<b>Eggs (% wet weight basis)</b>					
Moisture (%)	$58.39 \pm 0.50$	$58.22 \pm 0.32$	$58.46 \pm 0.29$	$58.24 \pm 0.34$	$58.65 \pm 0.11$
Protein (%)	$26.49 \pm 0.68$	$26.69 \pm 0.48$	$26.83 \pm 0.58$	$26.45 \pm 0.54$	$25.88 \pm 0.36$
Lipid (%)	$13.81 \pm 0.44$	$13.94 \pm 0.18$	$14.01 \pm 0.35$	$14.24 \pm 0.38$	$14.26 \pm 0.30$
<b>Muscle tissue (% dry weight basis)</b>					
Moisture (%)	$74.10 \pm 1.11$	$75.23 \pm 1.23$	$74.23 \pm 0.93$	$75.25 \pm 1.12$	$74.10 \pm 1.21$
Protein (%)	$64.49 \pm 0.27$	$62.74 \pm 0.28$	$62.84 \pm 0.78$	$62.73 \pm 0.59$	$62.58 \pm 0.50$
Lipid (%)	$31.60 \pm 0.13$	$32.35 \pm 0.33$	$32.62 \pm 0.46$	$32.64 \pm 0.48$	$32.59 \pm 0.45$
Ash (%)	$2.68 \pm 0.08$	$2.54 \pm 0.07$	$2.67 \pm 0.08$	$2.52 \pm 0.04$	$2.53 \pm 0.10$

Mean values in similar row with different letters are significantly different (Tukey's HSD test,  $P < 0.05$ ).

T<sub>1</sub>=100% FF; T<sub>2</sub>=75% FF + 25% LF; T<sub>3</sub>=50% FF + 50% LF; T<sub>4</sub>=25% FF + 75% LF; T<sub>5</sub>=100% LF.

**Fatty acid profile of muscle tissue.** The fatty acids of muscle tissue of *P. scalare* fed on different levels of formulated feed and live food is given in Table 5.



Table 5. Classification of fatty acid of total lipid in muscles of *P. scalare* broodstock fed on formulated feed and live food.

Fatty acid	T1	T2	T3	T4	T5
<b>Saturates</b>					
12:0	8.11±0.13 <sup>a</sup>	8.45±0.23 <sup>a</sup>	4.34±0.34 <sup>b</sup>	2.65±0.26 <sup>c</sup>	2.57±0.29 <sup>c</sup>
14:0	9.21±0.35 <sup>c</sup>	9.14±0.47 <sup>c</sup>	4.88±0.12 <sup>b</sup>	2.90±0.34 <sup>a</sup>	2.81±0.11 <sup>a</sup>
16:0	5.57±0.30 <sup>a</sup>	5.70±0.52 <sup>a</sup>	10.44±0.45 <sup>b</sup>	19.99±0.68 <sup>c</sup>	20.99±0.49 <sup>c</sup>
18:0	1.83±0.14 <sup>a</sup>	1.90±0.12 <sup>a</sup>	6.90±0.36 <sup>b</sup>	7.36±0.23 <sup>b</sup>	7.88±0.42 <sup>b</sup>
20:0	6.25±0.24 <sup>a</sup>	5.89±0.20 <sup>a</sup>	5.22±0.25 <sup>b</sup>	3.06±0.21 <sup>c</sup>	2.52±0.18 <sup>c</sup>
<b>Monounsaturates</b>					
18:1n-9	4.70±0.21 <sup>a</sup>	5.09±0.15 <sup>a</sup>	19.09±0.33 <sup>b</sup>	23.48±0.43 <sup>c</sup>	24.10±0.37 <sup>c</sup>
<b>n-6 Polyunsaturates</b>					
18:2n-6	12.20±0.28	11.84±0.33	11.81±0.26	11.66±0.23	11.42±0.19
<b>n-3 Polyunsaturates</b>					
18:3n-3	1.75±0.19 <sup>a</sup>	1.65±0.08 <sup>a</sup>	1.92±0.07 <sup>a</sup>	2.34±0.04 <sup>b</sup>	2.78±0.10 <sup>b</sup>
20:5n-3	12.31±0.27 <sup>c</sup>	11.78±0.24 <sup>c</sup>	4.42±0.16 <sup>b</sup>	2.30±0.30 <sup>a</sup>	2.26±0.13 <sup>a</sup>
22:6n-3	10.80±0.25 <sup>a</sup>	10.51±0.35 <sup>a</sup>	10.41±0.35 <sup>a</sup>	11.85±0.26 <sup>b</sup>	12.02±0.22 <sup>b</sup>

Mean values in similar row with different letters are significantly different (Tukey's HSD test,  $P < 0.05$ ).

### Discussion

Various studies have been conducted on the effect of nutrition on growth or reproduction of ornamental fish (Degani and Gur, 1992; Degani and Yehuda, 1996; Kruger et al., 2001; James and Sampath, 2002, 2004; Chong et al., 2004; Ling et al., 2006) but research into the impact of the combined feeding of live food and formulated feed on the reproductive performance of *P. scalare* is rare. Hence, this study was undertaken to examine the effects of feed quality on reproductive performance in marble angelfish, *P. scalare*.

From the different combinations of formulated feed and live food used, the highest relative fecundity and hatching rate was observed in fish fed on live food (mosquito larvae) indicating that the fatty acid composition of the diet influenced the reproductive performance of *P. scalare*. The best reproductive performance was observed in mosquito larvae fed groups (T4 and T5). This could be due to larger amounts of EPA (3.85%) and DHA (3.35%) than in the formulated feeds which contained 3.0% and 0.3% respectively. The balanced composition of lipids and essential n-3 and n-6 fatty acids in the live food might have stimulated gonad development and enhanced relative fecundity and hatching percentage in the angelfish. Similar observations were made for *Betta splendens*, where a mixed diet (50-100% substitution of live food) enhanced reproductive performance of fish (James and Sampath, 2002; 2003). In another study there was no significant difference in the number of eggs spawned in relation to diets (Luquet and Watanabe, 1986).

Type of diet significantly affected the percentage of hatching, and duration between spawns, but not the number of eggs per spawn in *P. scalare* (Degani and Yehuda, 1996). The short reproductive cycle of asynchronous fish might result in continuous oogenesis in adult females, therefore availability of the right type of diet is very important (Degani and Yehuda, 1996). During oogenesis, the oocytes synthesize vitellogenin (Degani et al., 1997) and in teleosts, oogenesis and spawning are regulated by the gonadotropin hormone (GtH). It is possible that live feed may contain some microelements essential for the secretion of GtHs. Another possibility is that the diet affects steroid production which is also important for oogenesis and the reproductive cycle (Degani and Boker, 1992).

Despite the relatively small quantity of live feed given in T3 diet, there was a significant improvement in the hatching percent when compared to fish fed only formulated feed. No benefits were observed in relative fecundity, inter spawning interval, hatching percent, and relative hatchling number when formulated feed was replaced with 25% live food. In our study presence or absence of mosquito larvae was more important

than the relative proportion of protein and lipid. This is supported by other reported observations (Degani and Yehuda, 1996).

Mosquito larvae may contain enzymes that enhance fish appetite, aid fish digestion, feed intake, growth, and reproductive performance. The developing embryo depends on nutrients present in egg yolk. The large size, active movement, and high protein (49.88%), and lipid (14.15%) content of mosquito larvae could have a stimulatory effect on the predatory responses of *P. scalare* which enhances relative fecundity, hatching rate, egg, and larvae quality. These results are supported by other studies for *P. scalare* (Degani and Yehuda, 1996; Ortega-Salas et al., 2009).

There was a 30% increase in fry of *Xiphophorus helleri* fed daily with a commercial diet supplemented with live *Daphnia* spp. (Kruger et al., 2001). This corresponds with our results. However in another study flake feed for maximum fry production of zebrafish (*Danio rerio*) was recommended (Markovich et al., 2007). This indicates that different fish species may have specific nutritional requirements for reproduction.

In the present study EPA and DHA played an important role in the reproductive performance of *P. scalare*. This is may be due to readily available n-3 HUFA for ovary development. Neural tissue, especially the eyes, forms a relatively large proportion of the total body mass of embryonic fish requirements, 22:6n-3, therefore any deficiency of this component during embryogenesis and larval development can lead to poor hatching rate. Another important function of (n-3) PUFA is their ability to produce eicosanoids, which are important in stress reaction and adaptation (Sargent et al., 1990).

In the present study, reproductive performance in terms of relative fecundity and hatching percentage of T3 was markedly improved compared to T1 and T2 but was lower than the groups fed with mosquito larvae T5 (100%) and T4 (75%) although there was no significant difference between the groups T3, T4 and T5 for hatching percentage and inter spawning interval. This improvement could possibly be due to the more balanced supply of both DHA and EPA. In addition, the amount of essential fatty acids in eggs correlates with inter-spawning interval, hatching rate, egg, and larvae quality (Blaxter, 1969).

Our study revealed a strong correlation between 16:00 and 18:1n-9 fatty acids with relative fecundity and hatching performance (hatching percentage and relative hatchling number) the reason being that that saturated and monounsaturated fatty acids are quantitatively more important than (n-3) PUFA in providing metabolic energy for egg development (Sargent, 1995).

Parameters used in assessing egg quality also include egg and oil globule diameter (Bromage et al., 1992). A number of environmental factors as well as size of brooders have been shown to influence fecundity and egg size. As fish size increases so does fecundity and diameter of eggs. The present study showed that in *P. scalare* fed with T4 and T5 diets, the egg size, and oil globule diameter was significantly larger than T1, T2, and T3. Egg size is an indicator of improved viability of the larvae (Blaxter, 1988). The difference in the size of the oil globule (table 3) did not have a significant effect on the hatching rate of the eggs.

The combined effect of formulated feed and live food significantly influenced the reproductive and hatching performance as well as egg and muscle composition of *P. scalare*. No significant differences were found for body protein, fat, and ash content among the treatments. It can be concluded that up to 50% live feed can be replaced with formulated feed for *P. scalare*, without any adverse effect on the reproductive performance of fish.

### Acknowledgements

I wish to thank Dr. Balasaheb Sawant Konkan Krishi Vidyapeeth and the relevant authorities in Ratnagiri, Maharashtra, India for providing me all the necessary facilities at the College of Fisheries, Ratnagiri.

### References

- AOAC**, 2006. Official Methods of Analysis, 18<sup>th</sup> ed. Association of Official Analytical Chemist, Washington, D.C. 1094 pp.
- Blaxter J.H.S.**, 1969. Development: eggs and larvae. In W.S. Hoar and D.J. Randall (eds), *Fish Physiology*, vol. 3, pp. 177-252. Academic Press, New York.
- Blaxter J.H.S.**, 1988. Pattern and variety in Development. In W.S. Hoar and D.J. Randall (eds), *Fish Physiology*, vol. XI A, pp. 1-58. Academic Press Inc., London, New York.
- Bromage N., Jones J., Randall C., Thrush M., Davies B., Springate J., Duston J. and G. Barker**, 1992. Broodstock management, fecundity, egg quality and timing of egg production in the rainbow trout (*Oncorhynchus mykiss*). *Aquaculture*, 100: 141-166.
- Chong A.S.C., Ishak S., Osman Z. And R. Hashim**, 2004. Effect of dietary protein levels on reproductive performance of female viviparous ornamental fish, swordtail *Xiphophorus helleri* (Poeciliidae). *Aquaculture*, 234:381-392.
- Degani G. and N. Gur**, 1992. Growth of juvenile *Trichogaster leerii* on diets with various protein levels. *Aquacult. Fish. Manag.*, 23:161-166.
- Degani G. and R. Boker**, 1992. Sensitivity to maturation inducing steroids and gonadotropin in the oocytes of blue gourami *Trichogaster trichopterus* (Anabantidae, Pallas, 1770). *Gen. Comp. Endocrinol.*, 85:430-439.
- Degani G. and Y. Yehuda**, 1996. Effects of diets on reproduction of angel fish *Pterophyllum scalare* (Cichidae). *Indian. J. Fish.*, 43:121-126.
- Degani G., Boker R., Gal E. and K. Jackson**, 1997. Oogenesis and steroid profiles during the reproductive cycle of female angelfish *Pterophyllum scalare* (Cichlidae). *Indian J. Fish.*, 44 (1):1-10.
- FAO**, 2010. Fish tat plus: Universal software for fishery statistical time species. Version 2.30 Fisheries Department, Fishery Information, Data and Statistics Unit Rome.
- Folch J., Lees M. and G.H. Sloane-Stanley**, 1957. A simple method for the isolation and purification of total lipids from animal tissue. *J. Biol.Chem.*, 226:497-509.
- Gunasekera R.M., Shim K.F. and T.J. Lam**, 1995. Effect of dietary protein level on puberty, oocyte growth and egg chemical composition in the tilapia, *Oreochromis niloticus* (L.). *Aquaculture*, 134:169-183.
- James R. and K. Sampath**, 2002. Effect of feed types on growth and fecundity in ornamental fish, *Betta splendens* (Regan). *Indian J. Fish.*, 49:279-285
- James R. and K. Sampath**, 2004. Effect of feed type on growth and fertility were studied in ornamental fish, *Xiphophorus helleri*, [\*Isr. J. Aquacult. - Bamidgeh\*](#), 56(4):264-273.
- James R. and K. Sampath**, 2003. Effect of animal and plant protein diets on growth and fecundity in ornamental fish. *Betta splendens* (Regan). [\*Isr. J. Aquacult. - Bamidgeh\*](#), 55(1):39-52.
- Kruger D.P., Britz P.J. and J. Sales**, 2001. The influence of livefeed supplementation on Growth and reproductive performance of swordtail (*Xiphophorus helleri* heckel 1848) broodstock. *Aqua. Sci. Cons.*, 3:265-273.
- Ling S., Hashim R., Kolkovski S. and A.S.C. Chong**, 2006. Effect of varying dietary lipid and protein levels on growth and reproductive performance of female swordtails *Xiphophorous helleri* (Poeciliidae). *Aquacult. Res.*, 37:1267-1275.
- Lowry O.H., Riousebrouch N.J., Far A.L. and R.J. Randall**, 1951. Protein measurement with the Folin reagent. *J. Biol. Chem.*, 193: 265-275.
- Luquet, P. and T. Watanabe**, 1986. Interaction: "Nutrition-reproduction" in fish. *Fish Physiol. Biochem.*, 2: 121-129.
- Mandal S.C., Sahu N.P., Kohli M.P., Das P., Gupta S.K. and S. Munilkumar**, 2010. Replacement of live feed by formulated feed: effect on the growth and spawning performance of Siamese fighting fish (*Betta splendens*, Regan, 1910). *Aquacult. Res.*, 41 (11): 1707-1716.
- Markovich M.L., Rizzuto N.V. and P.B. Brown**, 2007. Diet affects spawning in Zebrafish. *Zebrafish*, 4 (1): 69-74.



- Ortega-Salas A.A., Cortés G.I. and H. Reyes-Bustamante,** 2009. Fecundity, growth, and survival of the angelfish *Pterophyllum scalare* (Perciformes: Cichlidae) under laboratory conditions. *Rev. Biol. Trop.*, 57 (3):741-747.
- Rohlf F.J.,** 2004. tpsdig2 Stony Brook, NY: Department of Ecology and Evolution, State University of New York at Stony Brook.
- Sargent J.R.,** 1995. Origins and functions of egg lipids: nutritional implications. In: Bromage, N.R., Robert, R.J. (Eds.), Broodstock management and egg and larval quality. Blackwell, Oxford, pp. 353-372.
- Sargent J.R., Bell M.V., Henderson R.J. and D.R. Tocher,** 1990. In "Comparative Physiology, Vol. 5. Animal Nutrition and Transport Processes: Nutrition in Wild and Domestic Animals" ( J. Mellinger, J. P. Truchot, and B. Lahlou, eds.), p. 11. Karger, Basel.
- Schulz C., Bohm M., Wirth M. and B. Rennert,** 2007. Effect of dietary protein on growth, feed conversion, body composition and survival of pike perch fingerlings (*Sander lucioperca*). *Aquacult. Nutr.*, 13:373-380.
- Snedecor G.W. and W.G. Cochran,** 1967. Statistical methods, 6<sup>th</sup> ed. Oxford and IBH publishing Co., New Delhi, pp. 593.
- Zar J.H.,** 2005. Biostatistical Analysis, Fourth edition, Tan prints (I), Delhi, India, pp. 663.